

## **Nurse PrEP documentation template**

### **Assessment comments:**

See Progress note

### **Counseling/Follow-up**

- Client asymptomatic for acute HIV infection (no flu, mono – like illness in last 6 weeks).
- Discussed known safety risks with use of Truvada for a PrEP indication.
- Counseled on the importance of scheduled follow-up every 3 months, including regular HIV-1 screening tests (at least every 3 months), while taking Truvada for a PrEP indication.
- Instructed client to contact provider if they experience any of the following symptoms: fever, rash, joint pain, oral ulcers, fatigue, night sweats, sore throat, malaise, pain in muscles, loss of appetite.
- Counseled on the importance of adherence to daily dosing schedule and that effectiveness of medication is dependent on adherence to daily regimen.
- Counseled that Truvada for a PrEP indication should be used only as part of a comprehensive prevention strategy.
- Educated on practicing safer sex consistently and using condoms correctly.
- Discussed the importance of the individual knowing their HIV-1 status and, if possible, that of their partner(s).
- Discussed the importance of laboratory screening for sexually transmitted infections that can facilitate HIV-1 transmission
- Counseled on need for routine STI screenings Q 6 months.
- Reviewed HBV vaccination status. Ordered Hepatitis panel.
- Provided education on where information about Truvada for a PrEP indication can be accessed.
- Discussed potential adverse reactions and common side effects of Truvada medication.
- Reviewed the Truvada Medication Guide with the uninfected individual at high risk, copy given to client.
- HIV 4<sup>th</sup> generation, RPR, GC/chlamydia, Hepatitis panel, and serum creatinine collected and/or sent as PSC order for laboratory testing
- Client given condoms

### **For Women:**

- Reviewed reproductive life plan.
- Desires contraception, referred to Family Planning clinic.
- Pregnancy test negative